

# CHROMOSOME COUNTS IN ARUNDINOIDEAE, CHLORIDOIDEAE, AND POOIDEAE (POACEAE) FROM PAKISTAN<sup>1</sup>

M. Moinuddin,<sup>2</sup> Ahsan A. Vahidy,<sup>3</sup>  
and S. I. Ali<sup>2</sup>

## ABSTRACT

A total of 85 chromosome counts representing 45 species belonging to 29 genera in three subfamilies of Poaceae, viz., Arundinoideae, Chloridoideae, and Pooideae from Pakistan, are reported. The chromosome counts for six species, viz.: *Leptothrium senegalense* ( $n = 10$ ); *Piptatherum gracile* ( $n = 12$ ); *Poa sinaica* ( $n = 7$ ); *Sporobolus arabicus* ( $n = 18$ ); *Sporobolus nervosus* ( $n = 18$ ); and *Stipagrostis plumosa* ( $n = 22$ ), are new determinations. Chromosome counts for an additional 34 species are new for the flora of Pakistan. New cytotypes were detected in *Aeluropus lagopoides* ( $n = 20$ ), *Avena barbata* ( $n = 21$ ), *Diplachne fusca* ( $n = 20$ ), and *Ochthochloa compressa* ( $n = 10$ ).

The Poaceae are one of the largest families of flowering plants, comprising 620 genera and 10,000 species. Its members are widely dispersed in all regions of the world, occupying almost every type of habitat, often dominating the vegetation in savannas, prairies and steppes (Willis, 1973). In Pakistan, the family is represented by 158 genera and 492 species (Cope, 1982). Up to now chromosome numbers are known for 39 species (i.e., 7.92%) from Pakistan (Baquar & Anjum, 1969; Baquar & Saeed, 1969; Faruqi et al., 1979). The present contribution records the chromosome numbers of 45 species belonging to 29 genera in three subfamilies.

## MATERIALS AND METHODS

For meiotic counts, young unopened inflorescences were fixed in Carnoy's solution (3 parts absolute alcohol:1 part glacial acetic acid) for at least 24 hr., then stored at  $-4^{\circ}\text{C}$ . Slides were prepared by ordinary squash technique using 1.0% acetocarmine or 1.8% aceto-orcein.

For mitotic counts, young and healthy root tips from germinating seeds were pretreated either with 0.002 M 8-hydroxyquinoline (for 4–6 hr.) or with cold water (at  $1-2^{\circ}\text{C}$  for 24 hr.), fixed in Carnoy's solution for 1 hr., hydrolysed in 1 N HCl for 6–12 min. at  $60^{\circ}\text{C}$ , and squashed in 1.8% aceto-orcein.

In most of the cases photomicrographs were taken from temporary preparations. Desired slides were made permanent in euparal or Canada balsam. Voucher specimens are deposited in Karachi University Herbarium (KUH).

## OBSERVATIONS AND RESULTS

Information regarding the counts made, along with the basic number for the genus, are presented in Table 1. Basic numbers of genera were calculated from the lowest known gametic number in the genus. Chromosome numbers for six species are new to science and counts for thirty-four species are new to the flora of Pakistan, as they were not found to be reported in IPCN (Fedorov, 1974; Goldblatt, 1981, 1984, 1985, 1988; Moore, 1973, 1974).

## DISCUSSION

In the present study most of the chromosome counts that are in conformity with earlier reports have not been commented upon.


In the tribe Aristideae, counts for *Stipagrostis plumosa* ( $n = 22$ , Fig. 1) are reported here for the first time.

In the tribe Aeluropodeae, *Aeluropus lagopoides* ( $n = 20$ , Fig. 20) is found to be tetraploid on the basis of  $x = 10$ . Diploid and pentaploid

<sup>1</sup> We thank M. Qaiser, Z. A. Razaq, and S. M. Nadeem Ahsan of the Department of Botany, University of Karachi, for their help in the identification of vouchers. We also thank A. Ghafoor, S. Omer, T. Ali, Z. A. Razaq, B. Jahan, and N. Ahsan of the Department of Botany, University of Karachi, for collecting cytological material. This paper is a part of the work supported by National Science Foundation grant INT 8510318.

<sup>2</sup> Department of Botany, University of Karachi, Karachi-75270, Pakistan.

<sup>3</sup> Department of Genetics, University of Karachi, Karachi-75270, Pakistan.



FIGURES 1-6. Pollen mother cell meiosis in members of Poaceae.—1. *Stipagrostis plumosa* (Ghafoor 2776), metaphase-I  $n = 22$ .—2. *Diplachne fusca* (Siddiqui 101), diakinesis  $n = 20$ .—3. *Ochthochloa compressa* (T. Ali 752), diakinesis  $n = 10$ .—4. *Sporobolus arabicus* (T. Ali 842), metaphase-I  $n = 18$ .—5. *Sporobolus nervosus* (Siddiqui 131), diakinesis  $n = 18$ .—6. *Leptothrium senegalens* (Ahsan 68), diakinesis  $n = 10$ .

individuals have previously been reported for this species (Sindhe, 1980; Baquar & Saeed, 1969).

In tribe Eragrostideae, chromosome numbers for ten species (belonging to 6 genera) were determined. In *Dactyloctenium aegyptium*, we found  $n = 20$  in two and  $n = 23$  in one accession. On the basis of proposed basic numbers ( $x = 10, 12$ )

it can be suggested that individuals with  $n = 20$  were tetraploid, whereas  $n = 23$  represents a hyponeuploid based on  $x = 12$ . Our observations ( $n = 10, 20$ ), in two different specimens of *D. scindicum*, indicate the presence of cytotypes in this species.

Larsen (1963) had reported  $2n = 20$  ( $2x$ ) for

TABLE 1. Chromosome numbers in Poaceae from Pakistan. K.U. = Karachi University. D. G. Khan = town of Dera Ghazi Khan.

Taxon	Chromosome number <i>n</i>	Basic number <i>x</i>	Voucher
Subfamily Arundinoideae			
Tribe Arundineae			
* <i>Schismus barbatus</i> (L.) Thell. (Fig. 13)	6	6	Chitral: <i>Ghafoor</i> 2693, 3071; Makran: <i>T. Ali</i> 834
Subfamily Chloridoideae			
Tribe Aristideae			
* <i>Aristida adscensionis</i> L.	11	11	Makran: <i>T. Ali</i> 945
* <i>Aristida funiculata</i> Trin. & Rupr. (Fig. 19)	11		K.U. Campus: <i>Siddiqui</i> 4
** <i>Stipagrostis plumosa</i> (L.) Munro ex T. Anderson (Fig. 1)	22	22	Chitral: <i>Ghafoor</i> 2776
Tribe Aeluropodeae			
<i>Aeluropus lagopoides</i> (L.) Trin. ex Thwaites (Fig. 20)	20	10	Thatta: <i>Siddiqui</i> 150, 119; K.U. Campus: <i>Moin.</i> 71
Tribe Eragrostideae			
<i>Dactyloctenium aegyptium</i> (L.) Willd.	23	10, 12	K.U. Campus: <i>Razaq</i> 126
* <i>Dactyloctenium scindicum</i> Boiss.	10		K.U. Campus: <i>Moin.</i> 46; Layyah: <i>Ghafoor</i> 3744 K.U. Campus: <i>Moin.</i> 72
<i>Dactyloctenium scindicum</i>	20		K.U. Campus: <i>Ahsan</i> 56
* <i>Dinebra retroflexa</i> (Vahl) Panzer	10	10	Khushab: <i>Ghafoor</i> 3836
* <i>Diplachne fusca</i> (L.) P. Beauv. ex Roem & Schult. (Fig. 2)	20	10	Thatta: <i>Siddiqui</i> 101
<i>Eleusine indica</i> (L.) Gaertn.	9	9	K.U. Campus: <i>Ahsan</i> 6; Swat: <i>Ghafoor</i> 3977; Islamabad: <i>Ghafoor</i> 3469a
<i>Eleusine indica</i>	18		Makran: <i>Omer</i> 2112
* <i>Eragrostis cilianensis</i> (All.) Lutati ex F. T. Hubbard	10	10	K.U. Campus: <i>Moin.</i> 42
* <i>Eragrostis japonica</i> (Thunb.) Trin.	10		Lahore: <i>Ghafoor</i> 4358
* <i>Eragrostis minor</i> Host	10		Hasilpur: <i>Ghafoor</i> 3583
<i>Eragrostis minor</i>	20		Sargodha: <i>T. Ali</i> 1719
<i>Eragrostis pilosa</i> (L.) P. Beauv.	20		Sakesar: <i>T. Ali</i> 1605
* <i>Ochthochloa compressa</i> (Forssk.) Hilu (Fig. 3)	10	10	Makran: <i>T. Ali</i> 752
<i>Ochthochloa compressa</i>	20		Makran: <i>Omer</i> 2166, <i>T. Ali</i> 710; Safari Park, Karachi: <i>Ahsan</i> 38; K.U. Campus: <i>Razaq</i> 144; D. G. Khan: <i>Ghafoor</i> 3608
Tribe Chlorideae			
<i>Chloris barbata</i> Sw.	20	10	K.U. Campus: <i>Siddiqui</i> 11; Sajawal: <i>Ahsan</i> 26
* <i>Cynodon dactylon</i> (L.) Pers. (Fig. 14)	9	9	K.U. Campus: <i>Siddiqui</i> 31, 76, 90, <i>Moin.</i> 12; Dir: <i>Ghafoor</i> 2491
* <i>Tetrapogon tenellus</i> (Koen. ex Roxb.) Chiov. (Fig. 21)	10	10	K.U. Campus: <i>Moin.</i> 53
* <i>Tetrapogon villosus</i> Desf.	10		D. G. Khan: <i>Ghafoor</i> 3699; Makran: <i>T. Ali</i> 969

TABLE 1. Continued.





Taxon	Chromosome number <i>n</i>	Basic number <i>x</i>	Voucher
Tribe Sporoboleae			
** <i>Sporobolus arabicus</i> Boiss. (Fig. 4)	18	9	Makran: <i>T. Ali</i> 842, 846; Cholistan: <i>Ghafoor</i> 3550
* <i>Sporobolus coromandelianus</i> (Retz.) Kunth	18		K.U. Campus: <i>Moin.</i> 56
* <i>Sporobolus diander</i> (Retz.) P. Beauv.	18		Pail: <i>Ghafoor</i> 3776
** <i>Sporobolus nervosus</i> Hochst. (Fig. 5)	18		Thatta: <i>Siddiqui</i> 131; Darsanochano: <i>Siddiqui</i> 80
Tribe Zoysieae			
** <i>Leptothrium senegalense</i> (Kunth) W. D. Clayton (Fig. 6)	10	10	K.U. Campus: <i>Ahsan</i> 68
* <i>Tragus berteronianus</i> Schult.	10	10	Hazara: <i>Omer</i> 2221; Mianwali; <i>T. Ali</i> 1856
* <i>Tragus roxburghii</i> Panigrahi	10		K.U. Campus: <i>Moin.</i> 55
Subfamily Pooideae			
Tribe Poeae			
* <i>Dactylis glomerata</i> L.	7	7	Chitral: <i>Ghafoor</i> 2450, 3286
<i>Dactylis glomerata</i>	14		Dir: <i>Ghafoor</i> 2421
* <i>Lolium rigidum</i> Gaudin (Fig. 7)	7	7	Chitral: <i>Ghafoor</i> 3274
* <i>Poa annua</i> L.	14	7	Chitral: <i>Ghafoor</i> 2430; Kashmir: <i>T. Ali</i> 573; Kalam: <i>Ghafoor</i> 3422
** <i>Poa sinaica</i> Steud. (Fig. 8)	7		Hazara: <i>Omer</i> 2720
Tribe Glycerieae			
* <i>Glyceria plicata</i> (Fries) Fries	20	5	Chitral: <i>Ghafoor</i> 2850
Tribe Aveneae			
* <i>Agrostis munroana</i> Aitch. & Hemsl. (Fig. 22)	21	7	Hazara: <i>Omer</i> 2778
* <i>Agrostis viridis</i> Gouan (Fig. 9)	21+2B		Dir: <i>Ghafoor</i> 2337
<i>Agrostis viridis</i>	14		Chitral: <i>Ghafoor</i> 2698, 2513, 3325, 2608
* <i>Alopecurus arundinaceus</i> Poir.	14	7	Hazara: <i>Omer</i> 2722
* <i>Avena barbata</i> Pott ex Link (Fig. 23)	21	7	Gilgit: <i>Omer</i> 2559
* <i>Phalaris minor</i> Retz. (Fig. 10)	14	7	K.U. Campus: <i>Jahan</i> 19
* <i>Polypogon monspeliensis</i> (L.) Desf. (Fig. 11)	14	7	Chitral: <i>Ghafoor</i> 2598, 3249; K.U. Campus: <i>Siddiqui</i> 34; Makran: <i>Omer</i> 2143
Tribe Stipeae			
** <i>Piptatherum gracile</i> Mez (Fig. 12)	12	12	Chitral: <i>Ghafoor</i> 3093
* <i>Piptatherum munroi</i> (Stapf) Mez	12		Gilgit: <i>Omer</i> 2464; Swat: <i>Ghafoor</i> 3412; Dir: <i>Ghafoor</i> 2407
Tribe Bromeae			
* <i>Bromus danthoniae</i> Trin. (Fig. 15)	7	7	Chitral: <i>Ghafoor</i> 2916
* <i>Bromus pectinatus</i> Thunb.	7		Gilgit: <i>Omer</i> 2426
* <i>Bromus tectorum</i> L. (Fig. 16)	7		Chitral: <i>Ghafoor</i> 2606


TABLE 1. Continued.

Taxon	Chromosome number <i>n</i>	Basic number <i>x</i>	Voucher
Tribe Triticeae			
* <i>Elymus dahuricus</i> Turcz. ex Griseb.	21	7	Gilgit: Omer 2570
* <i>Elymus semicostatus</i> (Nees ex Steud.) Melderis (Fig. 17)	14		Chitral: Ghafoor 2707, 2975
* <i>Hordeum vulgare</i> L. (Fig. 18)	2 <i>n</i> = 14	7	K.U. Campus: Moin. 31 (seed obtained from NARC, Islamabad)

\* Count new to flora of Pakistan.

\*\* Count reported for the first time.






FIGURES 13-18. Chromosomes in members of Poaceae.—13. *Schismus barbatus* (Ghafoor 2693), diakinesis  $n = 6$ .—14. *Cynodon dactylon* (Siddiqui 90), metaphase-I  $n = 9$ .—15. *Bromus danthoniae* (Ghafoor 2916), metaphase-I  $n = 7$ .—16. *Bromus tectorum* (Ghafoor 2606) diakinesis  $n = 7$ .—17. *Elymus semicostatus* (Ghafoor 2707), diakinesis  $n = 14$ .—18. *Hordeum vulgare* (Moin. 31), mitotic metaphase  $2n = 14$ .

*Diplachne fusca*, whereas our material showed  $n = 20$  (Fig. 2). This 4x cytotype in *D. fusca* is a new report.

In *Ochthochloa compressa* only tetraploids were

reported previously (Hiremath & Chennaveeraiah, 1982). One of the present reports of  $n = 10$  (diploid) for this species establishes a new ploidy level (Fig. 3).

FIGURES 7-12. Pollen mother cell meiosis in members of Poaceae.—7. *Lolium rigidum* (Ghafoor 3274), diakinesis  $n = 7$ .—8. *Poa sinaica* (Omer 2720), diakinesis  $n = 7$ .—9. *Agrostis viridis* (Ghafoor 2337), diakinesis. [Arrows indicate B-chromosomes.]  $n = 21 + 2B$ .—10. *Phalaris minor* (Jahan 19), diakinesis  $n = 14$ .—11. *Polypogon monspeliensis* (Ghafoor 2598), metaphase-II  $n = 14$ .—12. *Piptatherum gracile* (Ghafoor 3093), diakinesis  $n = 12$ .



FIGURES 19-23. Pollen mother cell meiosis in members of Poaceae.—19. *Aristida funiculata* (Siddiqui 4), diakinesis  $n = 11$ .—20. *Aeluropus lagopoides* (Siddiqui 150), diakinesis  $n = 20$ .—21. *Tetrapogon tenellus* (Moin. 53), metaphase-I  $n = 10$ .—22. *Agrostis munroana* (Omer 2778), diakinesis  $n = 21$ .—23. *Avena barbata* (Omer 2559), anaphase-I  $n = 21$ .

In the genus *Tetrapogon*, chromosome numbers in two species were observed. The count for *Tetrapogon tenellus* is contrary to a previous report,  $n = 18$  (Singh & Godward, 1960), however our count agrees with basic number ( $x = 10$ ) for the genus (Table 1, Fig. 21).

Chromosome numbers of four species of *Sporobolus* were determined in the tribe Sporoboleae;

all of them have  $n = 18$ . The counts for *Sporobolus arabicus* (Table 1, Fig. 4) and *S. nervosus* (Table 1, Fig. 5) are reported as new to science.

In Zoysieae, chromosome numbers of three species belonging to two genera were determined. The count for *Leptothrium senegalense* (Table 1, Fig. 6) is reported for the first time.

In subfamily Pooideae, chromosome numbers

for 19 species (belonging to 13 genera) in six tribes were determined. The count for *Poa sinaica* (Table 1, Fig. 8) is reported for the first time.

Chromosome numbers for six species belonging to five genera were determined in Aveneae. In *Agrostis viridis*, we observed 21 bivalents and two B-chromosomes (Fig. 9). A hexaploid cytotype of *Avena barbata* ( $n = 21$ ) is reported for the first time (Fig. 23).

In the tribe Stipeae, the count for *Piptatherum gracile* (Table 1, Fig. 12) is reported for the first time.

#### LITERATURE CITED

- BAQUAR, S. R. & V. ANJUM. 1969. Chromosome behaviour in *Cenchrus* complex from southern West Pakistan. *Portugaliae Acta Biologica*. XI(3-4): 288-296.
- & M. SAEED. 1969. Chromosome studies and polyploidy analysis in grasses of West Pakistan I. *Caryologia* 22: 103-111.
- COPE, T. A. 1982. Poaceae. In: E. Nasir & S. I. Ali (editors), *Flora of W. Pakistan* 143: 1-678.
- FARUQI, S. A., H. B. QURAIH & N. HALAL. 1979. Chromosome numbers and some morphological characteristics of some Andropogoneae from Pakistan. *Cytologia* 44: 585-605.
- FEDOROV, A. A. (editor). 1974. *Chromosome Number of Flowering Plants*. Otto Koeltz Science Publishers, West Germany (reprinted).
- GOLDBLATT, P. (editor). 1981. *Index to Plant Chromosome Numbers 1975-1978*. *Monogr. Syst. Bot. Missouri Bot. Gard.* 5: 1-553.
- . 1984. *Index to Plant Chromosome Numbers 1979-1981*. *Monogr. Syst. Bot. Missouri Bot. Gard.* 8: 1-427.
- . 1985. *Index to Plant Chromosome Numbers 1982-1983*. *Monogr. Syst. Bot. Missouri Bot. Gard.* 13: 1-224.
- . 1988. *Index to Plant Chromosome Numbers 1984-1985*. *Monogr. Syst. Bot. Missouri Bot. Gard.* 23: 1-264.
- HIREMATH, S. C. & M. S. CHENNAVEERAIHAH. 1982. Cytogenetical studies in the wild and cultivated species of *Eleusine* (Gramineae). *Caryologia* 35: 57-69.
- LARSEN, K. 1963. *Studies in the Flora of Thailand*. 14. Cytological studies in vascular plants of Thailand. *Dansk Bot. Ark.* 20(3): 211-275.
- MOORE, R. J. (editor). 1973. *Index to Plant Chromosome Numbers for 1967-1971*. *Regnum Vegetabile* 90: 1-539.
- . 1974. *Index to Plant Chromosome Numbers for 1972*. *Regnum Vegetabile* 91: 1-108.
- SINDHE, A. N. R. 1980. Chromosome Number Reports LXVIII. *Taxon* 29: 545.
- SINGH, D. N. & M. B. E. GODWARD. 1960. Cytological studies in the Graminae. *Heredity* 15(2-3): 193-199.
- WILLIS, J. C. 1973. *A Dictionary of the Flowering Plants and Ferns*. Cambridge Univ. Press, Cambridge. (8th edition revised by Airy Shaw.)